

First projection [nl34]

Rewrite relaxation function from [nl31] with projection operators from [nl33] and apply Dyson identity $(X + Y)^{-1} = X^{-1} - X^{-1}Y(X + Y)^{-1}$ ¹

$$\begin{aligned}\langle f_0|f_0\rangle c_0(z) &= \left\langle f_0 \left| \frac{1}{z + \imath L} \right| f_0 \right\rangle = \left\langle f_0 \left| \frac{1}{z + \imath LP_0 + \imath LQ_0} \right| f_0 \right\rangle \\ &= \left\langle f_0 \left| \frac{1}{z + \imath LQ_0} \right| f_0 \right\rangle - \left\langle f_0 \left| \frac{1}{z + \imath LQ_0} \imath LP_0 \frac{1}{z + \imath L} \right| f_0 \right\rangle.\end{aligned}$$

Simplify both terms:

- $\left\langle f_0 \left| \frac{1}{z} \left[1 + \frac{(-\imath)}{z} LQ_0 + \frac{(-\imath)^2}{z^2} LQ_0 LQ_0 + \dots \right] \right| f_0 \right\rangle = \frac{1}{z} \langle f_0|f_0\rangle.$
- $\left\langle f_0 \left| \frac{1}{z + \imath LQ_0} \imath L \right| f_0 \right\rangle \frac{1}{\langle f_0|f_0\rangle} \left\langle f_0 \left| \frac{1}{z + \imath L} \right| f_0 \right\rangle = \left\langle f_0 \left| \frac{1}{z + \imath LQ_0} \imath L \right| f_0 \right\rangle c_0(z).$

$$\Rightarrow c_0(z) = \frac{1}{z + \frac{1}{\langle f_0|f_0\rangle} \left\langle f_0 \left| \frac{z}{z + \imath LQ_0} \imath L \right| f_0 \right\rangle},$$

$$\begin{aligned}\left\langle f_0 \left| \frac{z}{z + \imath LQ_0} \imath L \right| f_0 \right\rangle &= \left\langle f_0 \left| \left[1 - (z + \imath LQ_0) \frac{1}{z + \imath LQ_0} + \frac{z}{z + \imath LQ_0} \right] \imath L \right| f_0 \right\rangle \\ &= \left\langle f_0 \left| (-\imath) LQ_0 \frac{1}{z + \imath LQ_0} \imath L \right| f_0 \right\rangle = \left\langle f_0 \left| (-\imath) LQ_0 \frac{1}{z + \imath Q_0 LQ_0} Q_0 \imath L \right| f_0 \right\rangle \\ &= \left\langle f_1 \left| \frac{1}{z + \imath L_1} \right| f_1 \right\rangle; \quad |f_1\rangle = Q_0|f_0\rangle = Q_0 \imath L|f_0\rangle, \quad L_1 = Q_0 LQ_0.\end{aligned}$$

Relaxation function after first projection expressed via memory function:

$$c_0(z) = \frac{1}{z + \Sigma_1(z)}, \quad \Sigma_1(z) = \frac{1}{\langle f_0|f_0\rangle} \left\langle f_1 \left| \frac{1}{z + \imath L_1} \right| f_1 \right\rangle.$$

Memory function $\Sigma_1(z)$ of original problem, $\{L, |f_0\rangle\}$, can be reinterpreted as the (as yet non-normalized) relaxation function of a new dynamical problem, $\{L_1, |f_1\rangle\}$.

Projection operator Q_0 acts as filter on the Liouvillian L , absorbing that part of dynamics dealt with explicitly in first projection. Explicit information contained in normalization constant of $\Sigma_1(z)$.

¹Direct consequence of operator identity $(X+Y)(X+Y)^{-1} = X(X+Y)^{-1} + Y(X+Y)^{-1} = 1$, here with $X \doteq z + \imath LQ_0$, $Y \doteq \imath LP_0$.